



Case Report

Vaginal leiomyoma in a post menopausal woman

Surya Jayaram^{1,*}, Sarojini Mayadevi¹

¹Dept. of Obstetrics and Gynecology, Women's Health, Aster Medcity, Ernakulam, Kerala, India



ARTICLE INFO

Article history:

Received 07-09-2020

Accepted 14-10-2020

Available online 13-03-2021

Keywords:

Vaginal

Leiomyoma

Postmenopausal

ABSTRACT

Leiomyomas or fibroids are the commonest gynaecological tumours. It is only very rarely reported as vaginal lesion. It is usually treated by excision of the lesion. This is an interesting case of a 72-year-old lady who presented with vaginal pain and found to have a firm, mobile mass of 4cms x 5cms size located in the middle third of anterior wall of vagina. MRI showed a well-encapsulated solid, round mass with peripheral enhancement localised to anterior vaginal wall. This mass was removed vaginally under anaesthesia. Histopathologic examination of the mass showed leiomyoma.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Leiomyomas or fibroids arising from uterus are the most common gynaecological tumours that are seen in about 20-40% of women after 35 years of age.¹ They are also seen in extrauterine sites and may be mistaken for benign lesions like urethral diverticulum, cystocele, abscesses or a vaginal malignancy.^{2,3} They are usually seen in women between 35 to 50 years of age.² Pre-operative diagnosis is usually difficult with imaging techniques like ultrasound and MRI. Diagnosis can be confirmed by postoperative histopathology.

2. Case Presentation

A 76 years old post-menopausal lady, para 2, presented to us with complaints of vaginal pain for past 2 weeks. She had no history of discharge per vaginam, bleeding per vaginam or urinary symptoms. She had attained menopause at 46 years of age. She gave history of asymptomatic small uterine fibroids in ultrasound scans done in her pre-menopausal period. There was history of carcinoma breasts and carcinoma endometrium in her mother.

On pelvic examination she had a firm, smooth, non-tender mass of 5x4cms arising from anterior vaginal wall in mid vaginal region. Uterus and cervix were found atrophied.

2.1. Investigations

Ultrasound scan and MRI was done which showed a well-defined solid round, mass of 4.5x5.5cms size arising from middle third of anterior vaginal wall in submucosal plane with peripheral enhancement, likely to be leiomyoma.

2.2. Differential diagnosis

Papilloma, haemangioma, mucous polyp, leiomyoma from anterior vaginal wall.

2.3. Treatment

The tumour was enucleated vaginally through a transverse incision over anterior vaginal wall. Urethra was protected during excision by placing a urethral Foleys catheter. The tumour was enucleated by blunt dissection. Vaginal wall closed by interrupted sutures. The tumour was then sent for histopathological examination which showed a 4 × 5 cm solid mass with a typical whorling pattern in the cut section (Figures 1 and 2). On microscopic examination

* Corresponding author.

E-mail address: suryaj1@yahoo.co.in (S. Jayaram).

features were consistent with leiomyoma - spindle shaped cells organized in bundles and fascicles with vascularized connective tissue in between. Individual cells were seen to have moderate amount of eosinophilic cytoplasm, cigar shaped nucleus with sparse mitosis. (Figure 3)

Patient was followed up at 2 weeks, 6 weeks and 1 year and was symptom free.



Fig. 1: Gross appearance of vaginal leiomyoma



Fig. 2: Cross section of leiomyoma showing whorled pattern

3. Discussion

Vaginal leiomyomas are uncommon with only about 300 cases reported so far.² Denis de Leyden reported the first case of vaginal leiomyoma back in 1733.² Bennett and Erlich had reported that at Johns Hopkins Hospital they found only nine cases in 50,000 surgical specimens examined and only one case in 15,000 autopsies reviewed.³ Leiomyomas from vaginal wall are extremely rare -




Fig. 3: Microscopic appearance of vaginal leiomyoma; Left: Spindle cells arranged in fascicles and bundles separated by vascularized connective tissue. Right: Individual cells have moderate amount of eosinophilic cytoplasm, cigar shaped nucleus with sparse mitosis

constituting only 4.5% of solid tumours of vagina.⁴

It has been found that leiomyomas occurring in vaginal wall are usually seen in Caucasian women aged between 35 and 50 years.³ They are most commonly seen as solitary well-defined lesions arising from the midline area of anterior vaginal wall.¹ Depending on location and size of the lesion patients may present with a vaginal, vaginal bleeding, lower abdominal pain, urinary symptoms due to urinary bladder compression /obstruction or dyspareunia.^{5–8}

Transvaginal ultrasound is the recommended first-line imaging modality for vaginal masses as per American College of Radiology Appropriateness criteria.⁹ Ultrasound as first line diagnostic modality is limited by its operator dependence and possibility of bypassing the lesion in perineum or lower vagina when the ultrasound probe is introduced into vaginal fornix.⁶ MRI is recommended if transvaginal ultrasound imaging is non-diagnostic.¹⁰ In MRI myomas usually appear as rounded homogenous lesions with signal intensities similar to that of the myometrium, with multiple calcifications. Leiomyosarcomas are characterised by absence of calcifications in the lesion.¹¹ If intra-lesional haemorrhage is present it usually points towards a sarcomatous lesion.^{12,13}

The treatment of choice of vaginal leiomyoma is surgical excision either through vaginal or abdomino perineal approach with utmost care so as not to damage underlying structures like urinary bladder, urethra or rectum. Sometimes reconstructive surgery with simple flaps or myocutaneous flaps might be necessary to correct the defects after removal of large vaginal leiomyomas.^{4,14}

The gold standard for diagnosis of vaginal leiomyoma is histopathological examination. It has the additional benefit of ruling out any focus of malignancy. If malignancy is found, further treatment either in the form of repeat excision or chemotherapy/radiotherapy may be needed. The histological features of leiomyosarcoma include the presence of two of the following three histological features:

cytological atypia, coagulative tumour cell necrosis and 10 or more mitoses per 10 high-power microscopic fields. Now, the presence of coagulative tumour cell necrosis and cytological atypia are considered as the most important features in distinguishing a benign from a malignant smooth muscle tumour.¹⁵

3.1. Learning Points/take Home Messages

Vaginal leiomyomas are rare conditions which can be misdiagnosed easily. Even though leiomyomas are oestrogen dependent tumours which are more common in reproductive age group, they can be found in post-menopausal women also. Use of proper imaging modalities followed by careful excision of the lesion and its histopathological examination is the treatment of choice.

4. Source of Funding

None.

5. Conflict of Interest

The authors declare that there is no conflict of interest.

References

1. Thompson RJ, Rock JA. Te Linde's Operative Gynecology. Ed, editor. Philadelphia: Lippincott Williams & Wilkins; 1992.
2. Young SB, Rose PG, Reuter KL. Vaginal fibromyomata: two cases with preoperative assessment, resection, and reconstruction. *Obstet Gynecol.* 1991;78(5):972–4.
3. Bennett EM. Myomata of vagina. *Am J Obstet Gynecol.* 1941;42:314–20.
4. Singh R, Yadav P, Kaur H. Vaginal Leiomyoma: A Rare Presentation. *J South Asian Feder Obst Gynecol.* 2014;6(2):112–3.
5. Pulfus E, Newcomer J. Guest Editorial. *Obstet Gynecol Surv.* 1999;54(3):149–50. doi:10.1097/00006254-199903000-00001.
6. and A De ICI, Pati S. Vaginal leiomyoma. *J Midlife Health.* 2011;2(1):42–3.
7. Sim CH, Lee JH, Kwak JS, Song SH. Necrotising vaginal leiomyoma mimicking amalignant neoplasia. *Obstet Gynecol Sci.* 2014;57(6):560–3.
8. Imai A, Furui T, Hatano Y, Suzuki M, Suzuki N, Goshima S. Leiomyoma and Rhabdomyoma of the vagina. *J Obstet Gynecol.* 2008;28(6):563–6.
9. American College of Radiology," 25 Aug 2016. [Online]. Available from: http://www.acr.org/secondarymainmenucategories/quality_safety/app_criteria.aspx.
10. Walker DK, Salibian RA, Salibian AD, Belen KM, Palmer SL. Overlooked diseases of the vagina: a directed anatomic-pathologic approach for imaging assessment. *Radiographics.* 2011;31:1593–8.
11. Elsayes KM, Narra VR, Dillman JR, Velcheti V, Hameed O, Tongdee R, et al. Vaginal masses: magnetic resonance imaging features with pathologic correlation. *Acta Radiol.* 2007;48(8):921–33. doi:10.1080/02841850701552926.
12. Den CAMAV, Bosch T. Screening for uterine tumors. *Best Pract Res Clin Obstet Gynecol.* 2012;26:257–66.
13. Samuel A, Fennessy FM, Tempany CMC, Stewart EA. Avoiding treatment of leiomyosarcomas: the role of magnetic resonance in focused ultrasound surgery. *Fertil Steril.* 2008;90(3):850.e9–e12. doi:10.1016/j.fertnstert.2007.08.019.
14. Gowri R, Soundararaghavan S, Oumachigui A, Sistla SC, Iyengar KR. Leiomyoma of vagina-an unusual presentation. *J Obstet Gynaecol Res.* 2003;29(6):395–8.
15. Watanabe S. Uterine leiomyoma versus leiomyosarcoma : a new attempt a differential diagnosis based on cellular characteristics. *Histopathol.* 2006;48:563–8.

Author biography

Surya Jayaram, Senior Specialist

Sarojini Mayadevi, Senior Consultant

Cite this article: Jayaram S, Mayadevi S. Vaginal leiomyoma in a post menopausal woman. *Indian J Obstet Gynecol Res* 2021;8(1):130-132.