

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Obstetrics and Gynecology Research

Journal homepage: www.ijogr.org

Case Report

Ascites in preeclampsia: A case report

Sharmeen Vazifdar^{1,*}, Urmila Gavali¹

¹Dept. of Obstetrics and Gynecology, Dr Vithalrao Vikhe Patil Foundations Medical College, Ahmednagar, Maharashtra, India



ARTICLE INFO

Article history:

Received 27-01-2022

Accepted 09-02-2022

Available online 20-05-2022

Keywords:

Preeclampsia

Pregnancy

ABSTRACT

Preeclampsia complicates approximately 2-8% of all pregnancies and is an important cause of maternal morbidity and mortality. Ascites in preeclampsia is a rare complication. We here are presenting a case of a 28-year primigravida with 34 weeks of gestation, a known case of preeclampsia, that was referred to our institute from a primary health centre in view of impending eclampsia. Massive ascites is an unusual complication of preeclampsia which may lead to maternal respiratory compromise, which calls for active termination of the pregnancy within 24–48 hours as it cannot be cured by medical treatment. Its rarity may be due to under-reporting. The incorporation into clinical practice of evaluating the amount of ascites in patients with preeclampsia might prompt the obstetrician to more intensive and more frequent maternal and fetal surveillance to avoid maternal and fetal complications.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Pre-eclampsia is defined as new onset hypertension and proteinuria after 20 weeks of gestation.¹ Release of vasoconstrictive agents, endothelial damage, hyperpermeability of the capillaries and microangiopathic hemolysis constitute the basic pathophysiology of this condition.² Preeclampsia complicates approximately 2-8% of all pregnancies and is an important cause of maternal morbidity and mortality.³ Massive ascites in preeclampsia is a rare complication. It is a manifestation of the multisystem organ exaggeration of preeclampsia. The occurrence of ascites can lead to adverse maternal outcome.⁴ The rarity of this condition may be due to under-reporting. Here we are presenting a case of preeclampsia in which massive ascites happened to be one of the findings intraoperatively.

2. Case Report

A 28-year primigravida with 34 weeks of gestation was referred to our institute from a primary health centre in view of impending eclampsia. Patient complained of headache since morning with no other premonitory signs. She was a known case of preeclampsia and was on tablet labetalol 100 mg twice a day since the past 10 days. The examination showed bilateral pedal oedema. The patient's pulse was 98 per minute and blood pressure was 200/120mm of mercury. The patient's respiratory rate was 18 beats per minute however chest was auscultated and found to be clear. The per abdomen examination showed a 32 weeks sized uterus with cephalic presentation. Abdominal wall oedema was seen and fluid thrill could be elicited suggestive of ascites. The fetal heart rate was found to be 136 per minute. On per vaginal examination, cervical os was closed. On urine dipstick test, urine protein was +3. All routine laboratory investigations like complete blood count, liver and renal function tests were sent and subsequently the reports were within normal limits. The

* Corresponding author.

E-mail address: sharmeenvazifdar@yahoo.com (S. Vazifdar).

patient's PIH profile consisting of PT (17 seconds) INR (1.42), LDH (1140IU/L), serum uric acid (6.5mg/dl) and serum albumin(3g/dl) were deranged. NST was reactive. Patient was given 12mg betamethasone for fetal lung maturity. Magnesium sulphate prophylaxis was given as per Prichardt's regimen. Injection labetalol 20mg was administered intravenously after which her blood pressure was recorded as 180/110, that is, there was no significant drop in blood pressure. The dose was subsequently repeated. The patient was taken up for an emergency lower segment caesarean section. On laparotomy, approximately 1.5 litres straw coloured ascitic fluid was seen. A 1950gm male fetus was delivered with evidence of IUGR. Magnesium sulphate was continued for 24hrs after surgery. Mother and baby were both discharged on postoperative day 10. Ultrasonography done at time of discharge showed no ascites, urine albumin was negative and blood pressure was 130/88mm of mercury. Antihypertensives consisting of oral labetalol 100mg twice as day were continued and patient was asked to follow up after 2 weeks.

3. Discussion

Preeclampsia may have variable clinical presentation. The presence of reported ascites in the literature varies between 5% to 20%.⁵ Although the cause of ascites in preeclampsia is obscure, most probable explanation is generalized capillary leak due to endothelial cell dysfunction and reduced intravascular oncotic pressure.⁶ It is this low colloid osmotic pressure that results in effusion such as ascites.⁷ Approximately 83 cases of large volume ascites have been reported in the setting of preeclampsia.⁸ Many cases of preeclampsia with ascites have been recorded mainly in the form of case reports.^{8–11} In an 11-year study of 23 patients with pregnancy induced hypertension (PIH) and ascites, Cong and Wang found that the incidence of ascites was 21.6 in 1,000 in severe PIH, and that the presence of this condition warrants termination of pregnancy as it can not be cured using medical treatment.¹⁰ Vaijyanth et al estimated the incidence of ascites in preeclampsia as 1/1000 in their report.¹² In a study conducted in 2019 by Mbonyizina C et al. out of 106 pts with severe preeclampsia 46 had ascites on ultrasonography. There was increased morbidity seen in these patients.⁵ In many studies massive ascites was noted at the time of caesarean section.¹³

Exploration of the abdomen should be conducted when ascites is found intraoperatively, to rule out intraperitoneal tumors. The abdominal cavity was explored at the time of cesarean section in our patient, no demonstrable organic cause, benign or malignant, other than pre-eclampsia could be found to explain the ascites. The clinical course of this case indicated that the underlying cause of the massive ascites was preeclampsia. The amount of ascites decreased after hypertension and proteinuria resolved. After the classic clinical signs of pre-eclampsia (hypertension, proteinuria,

edema) disappeared, the amount of peritoneal fluid also normalized. Massive ascites is an unusual complication of preeclampsia. It may lead to maternal respiratory compromise, which calls for active termination of the pregnancy within 24–48 hours as it cannot be cured by medical treatment. Its rarity may be due to under-reporting.

4. Conclusion

Careful clinical and ultrasonographic examinations are necessary to detect massive ascites in pregnancy. The incorporation into clinical practice of evaluating the amount of ascites in patients with preeclampsia might prompt the obstetrician to more intensive and more frequent maternal and fetal surveillance to avoid maternal and fetal complications.

5. Source of Funding

None.

6. Conflict of Interest

None.

References

1. Phipps E, Prasanna D, Brima W, Jim B. Preeclampsia: updates in pathogenesis, definitions, and guidelines. *Clin J Am Soc Nephrol.* 2016;11(6):1102–13.
2. Boeldt DS, Bird IM. Vascular adaptation in pregnancy and endothelial dysfunction in preeclampsia. *J Endocrinol.* 2017;232(1):R27–R44.
3. Köseoglu SB, Deveer R, Camuzcuoglu A, Kasap B, Camuzcuoglu H. A rare postpartum complication of severe preeclampsia: Massive ascites and pleural effusion. *J Turk German Gynecol Assoc.* 2016;17:S170.
4. Alabrah PW, Agbo J, Agoro EY. Preeclampsia Complicated by Antenatal and Postnatal Ascitis. *Eur J Med Health Sci.* 2019;1(5):10–3. doi:10.24018/ejmed.2019.1.5.106.
5. Mbonyizina C, Ntirushwa D, Bazzett-Matabele L, Ntsumbumuyange D, Rulisa S, Magriples U. Point of care ultrasound: does the presence of ascites in severe pre-eclampsia correlate with poor maternal and neonatal outcome? *Trop Med Int Health.* 2019;24(8):1018–22.
6. Koseoglu SB, Deveer R, Camuzcuoglu A, Kasap B, Camuzcuoglu H. Massive Ascites and Pleural Effusion in Preeclampsia. *J Clin Diagn Res.* 2017;11(2):QD08.
7. Pradhan P, Sherpa K, Joshi A, Pathak S. Massive ascites in severe pre-eclampsia: a rare complication. *Nepal Med Coll J.* 2012;14(4):342–4.
8. Morton A. Severe pre-eclampsia complicated by multiple serous effusions. *Aust New Zealand J Obstet Gynaecol.* 2004;44(4):360–1.
9. Chawla S, Kumar P, Bhalla M. Case of ascites in pre-eclampsia. *Med J Armed Forces India.* 2012;68(3):257–9.
10. Ding DC, Hsu S, Su HY, Chu TW. Massive Ascites in Severe Pre-Eclampsia. *Taiwanese J Obstet Gynecol.* 2005;44(4):384–6.
11. Ko ML, Chen SC, Huang LW, Chang JZ, Huang LW, Hwang JL, et al. Massive ascites complicating pre-eclampsia. *Taiwanese J Obstet Gynecol.* 2005;44(3):267–9.
12. Vaijyanath AM, Nayar B, Malhotra N, Deka D. Massive ascites in severe pre-eclampsia: A rare complication. *J Obstet Gynaecol Res.* 2002;28(4):199–202.
13. Ghazali S, Al-Domyati M. Severe Hyponatremia and Ascites Associated with Preeclampsia. *Crescent J Med Biol Sci.* 2019;6(1):132–5.

Author biography

Sharmeen Vazifdar, Post Graduate Resident

Urmila Gavali, Professor

Cite this article: Vazifdar S, Gavali U. Ascites in preeclampsia: A case report. *Indian J Obstet Gynecol Res* 2022;9(2):296-298.